| Please check the examination details below before entering your candidate information | | | | | | | | |
|---|----------------------------|--|--|--|--|--|--|--|
| Candidate surname | Other names | | | | | | | |
| Pearson Edexcel Level 1/Level 2 GCSE (9–1) | re Number Candidate Number | | | | | | | |
| Wednesday 12 J | lune 2019 | | | | | | | |
| Morning (Time: 1 hour 45 minutes) | Paper Reference 1CH0/2H | | | | | | | |
| Chemistry Paper 2 | | | | | | | | |
| | Higher Tier | | | | | | | |
| You must have: Calculator, ruler | Total Marks | | | | | | | |

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided
 there may be more space than you need.
- Calculators may be used.
- Any diagrams may NOT be accurately drawn, unless otherwise indicated.
- You must show all your working out with your answer clearly identified at the end of your solution.

Information

- The total mark for this paper is 100.
- The marks for each question are shown in brackets
 use this as a guide as to how much time to spend on each question.
- In questions marked with an asterisk (*), marks will be awarded for your ability to structure your answer logically showing how the points that you make are related or follow on from each other where appropriate.
- A periodic table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



P56423A



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Answer ALL questions. Write your answers in the spaces provided.

Some questions must be answered with a cross in a box \boxtimes . If you change your mind about an answer, put a line through the box \boxtimes and then mark your new answer with a cross \boxtimes .

1 (a) (i) Titanium(IV) oxide is an ionic solid. Many ionic solids are soluble in water.

Titanium(IV) oxide is not soluble in water. Its other physical properties are typical of ionic solids.

Predict **one** other physical property of titanium(IV) oxide that would be typical of ionic solids.

(1)

(ii) The formula of titanium(IV) oxide is TiO₂.

Deduce the charge of the titanium ion in titanium(IV) oxide.

(1)

- (b) Nanoparticles are very small particles that have unusual properties.
 - (i) Particles less than 100 nanometres in size are classified as nanoparticles.

100 nanometres is

(1)

- \triangle **A** 1×10^{-4} metres
- \blacksquare **B** 1×10^{-5} metres
- \square **C** 1×10^{-7} metres
- \square **D** 1×10^{-9} metres

DO NOT WRITE IN THIS AREA

| (ii) Nanoparticles of titanium(IV) oxide are used in some sunscreens. | |
|---|----------------|
| Describe a reason why nanoparticles of titanium(IV) oxide are used in some su | unscree (2) |
| | |
| | |
| | |
| (iii) Some people are concerned that there is a risk when sunscreens containing nanoparticles are used. | |
| Explain a possible risk associated with using nanoparticles in sunscreens. | (2) |
| | |
| | |
| | |
| (Total for Question 1 = 7 mag | arks) |



DO NOT WRITE IN THIS AREA

- 2 Most of the fuels used today are obtained from crude oil.
 - (a) Which statement about crude oil is correct?

(1)

- ☑ A crude oil is a compound of different hydrocarbons
- **B** crude oil is a mixture of hydrocarbons
- C crude oil contains different hydrocarbons, all with the same molecular formula
- **D** crude oil is an unlimited supply of hydrocarbons
- (b) Crude oil is separated into several fractions by fractional distillation. Two of these fractions are kerosene and diesel oil.
 - (i) State a use for each of these fractions.

(2)

kerosene

diesel oil

(ii) Figure 1 shows where the fractions kerosene and diesel oil are produced in the fractionating column.




Figure 1

Kerosene is obtained higher up the column than diesel oil. Kerosene and diesel oil fractions have slightly different properties.

Choose a property.

State how this property for kerosene compares with the property for diesel oil.

(1)

| property | |
|----------|--|
| | |

comparison

(c) Figure 2 shows the formulae of a molecule of butane and of a molecule of pentane. Butane and pentane are neighbouring members of the same homologous series.




Figure 2

(i) Explain, using these formulae, why butane and pentane are neighbouring members of the same homologous series.

(2)

(ii) Butane has the formula C₄H₁₀.

Calculate the mass of carbon in 100 g of butane.

Give your answer to three significant figures.

(relative atomic masses: H = 1.00, C = 12.0; relative formula mass: $C_4H_{10} = 58.0$)

You must show your working.

(3)

(Total for Question 2 = 9 marks)

3 (a) An aluminium atom has the atomic number 13 and the mass number 27.

Which row shows the numbers of subatomic particles present in an aluminium ion, Al³⁺?

(1)

| | | protons | neutrons | electrons |
|---|---|---------|----------|-----------|
| X | A | 13 | 14 | 13 |
| X | В | 13 | 14 | 10 |
| X | C | 14 | 13 | 10 |
| X | D | 14 | 13 | 17 |

(b) Magnesium burns in excess oxygen to form magnesium oxide. The balanced equation for this reaction is

$$2Mg + O_2 \rightarrow 2MgO$$

Starting with 1.35g of magnesium, calculate the maximum mass of magnesium oxide that could be formed in this reaction. (relative atomic masses: O = 16.0, Mg = 24.0)

You must show your working.

(3)

mass of magnesium oxide =g

(c) Chlorine reacts with hydrogen to form hydrogen chloride.

Write the balanced equation for this reaction.

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(d) Sodium reacts with chlorine to form sodium chloride.

The electronic configuration of the sodium atom is 2.8.1 and the electronic configuration of the chlorine atom is 2.8.7.

Give the electronic configurations of the ions formed.

(2)

Na⁺

Cl⁻

(Total for Question 3 = 9 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

| | | | nce of yeast. | |
|-----|------|----|--|-----|
| | | | action is carried out at 30 °C. | |
| | | | n why the reaction is carried out at a temperature of 30°C rather than at a crature of 80°C. | (2) |
| | | | | |
| (b) | | | ol, C₂H₅OH, can be converted into ethanoic acid, CH₃COOH. | |
| | (i) | In | this reaction ethanol is | (1) |
| | X | A | hydrated | |
| | X | В | oxidised | |
| | X | | polymerised | |
| | × | D | reduced | |
| | (ii) | Dr | aw the structure of a molecule of ethanoic acid, CH₃COOH, showing all | |
| | | CO | valent bonds. | (2) |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |
| | | | | |

DO NOT WRITE IN THIS AREA

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(c) (i) The apparatus in Figure 3 can be used to investigate the temperature rise produced in a known mass of water when a sample of ethanol is burned.




Figure 3

The first steps of the method are

- 1. put 100cm³ of water into a beaker
- 2. determine the mass of the burner containing ethanol
- 3. measure the initial temperature of the water
- 4. place the burner under the beaker of water
- 5. light the wick

Describe the remaining steps of the method that are needed to determine the mass of ethanol required to raise the temperature of the water by 30 °C.

| | | |
|------|------|--|
| | | |
| | | |
| | | |
| | | |
| | | |


(ii) In a different experiment, separate samples of the alcohols methanol, ethanol, propanol, butanol and pentanol were burned to determine the mass of each alcohol that needs to be burned to raise the temperature of 100 cm³ water by 10 °C.

| alcohol | number of carbon atoms in one molecule of alcohol | mass of alcohol burned in g |
|----------|---|-----------------------------|
| methanol | 1 | 0.37 |
| ethanol | 2 | 0.28 |
| propanol | 3 | 0.25 |
| butanol | 4 | 0.23 |
| pentanol | 5 | 0.22 |

Draw a graph of the mass of each alcohol required to raise the temperature of 100cm³ of water by 10 °C against the number of carbon atoms in one molecule of that alcohol.

(3)

mass of alcohol burned in g



number of carbon atoms in one molecule of alcohol

(Total for Question 4 = 11 marks)

NOT WRITE IN THIS AREA

- **5** (a) Carbon dioxide is one of the gases in the Earth's atmosphere.

 The percentage of carbon dioxide in the Earth's atmosphere has changed over time.
 - (i) Which row of the table shows the approximate percentage of carbon dioxide thought to be in the Earth's early atmosphere and how this percentage changed to form the Earth's atmosphere today?

(1)

| | approximate percentage of carbon dioxide in the Earth's early atmosphere | change in percentage carbon dioxide to form the Earth's atmosphere today. | | | | |
|---|--|---|--|--|--|--|
| Α | 5 | increased | | | | |
| В | 5 | decreased | | | | |
| C | 95 | increased | | | | |
| D | 95 | decreased | | | | |

(ii) The actual percentage of carbon dioxide in the Earth's atmosphere today varies.

Explain **two** factors that cause the percentage of carbon dioxide in today's atmosphere to vary.

(4)

| ı | |
|---|----------|
| | |
| | |
| | |
| | factor 2 |
| | |
| | |
| | |

DO NOT WRITE IN THIS AREA

| Explain why carbon dioxide has a low boiling point. | (2) |
|--|-----|
| | |
| | |
| | |
| (c) Calculate the number of molecules in 0.11 g of carbon dioxide. Give your answer to two significant figures. | |
| (relative formula mass : $CO_2 = 44$ Avogadro constant = 6.02×10^{23}) | (3) |
| | |
| | |
| number of molecules = | |
| (Total for Question 5 = 10 mar | ks) |



- 6 Some of the elements in the periodic table are metals.
 - (a) The electronic configuration of a metal is 2.8.3

Which row shows the group and period of the periodic table where this metal is found?

1)

| | group | period |
|-----|-------|--------|
| ⊠ A | 2 | 3 |
| В | 2 | 8 |
| ⊠ C | 3 | 2 |
| □ D | 3 | 3 |
| | | |

- (b) Lithium, potassium and rubidium are alkali metals.
 - (i) Describe what you would see when a small piece of rubidium is dropped on to water.

(2)

(ii) The electronic configuration of lithium is 2.1 The electronic configuration of potassium is 2.8.8.1 Lithium is less reactive than potassium.

Explain, in terms of their electronic configurations, why lithium is less reactive than potassium.

(3)

| | |
|------|------|------|------|------|------|------|--|
| | | | | | | | |
| | |
| | |
| | |
| | | | | | | | |



DO NOT WRITE IN THIS AREA

| (c) Lithium has two naturally occurring isotopes, lithium-6 and lithium-7. | |
|--|-----|
| A sample of lithium contains 7.59% of lithium-6 92.41% of lithium-7. | |
| Calculate the relative atomic mass of lithium in this sample. | |
| Give your answer to two decimal places. You must show your working. | |
| Tou must show your working. | (4) |
| | |
| | |
| | |
| | |
| | |
| | |
| relative atomic mass of lithium | 1 = |
| (Total for Question 6 - | |



DO NOT WRITE IN THIS AREA

| 7 | Ethene, C ₂ H ₄ , is an unsaturated hydrocarbon. | |
|---|---|-----|
| | (a) Explain why ethene is an unsaturated hydrocarbon . | (2) |
| | | (2) |
| | | |
| | | |
| | | |
| | | |
| | (b) A sample of ethene is burned completely in oxygen. | |
| | Write the balanced equation for this reaction. | (3) |
| | | (3) |
| | / | |
| | (c) Ethene can be polymerised to form poly(ethene). | |
| | Describe what you would see when a sample of ethene and a sample of poly(ethene) are shaken with separate, small volumes of bromine water. | |
| | | (3) |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

DO NOT WRITE IN THIS AREA

| | molecular formula = |
|--|---------------------|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | (3) |
| (relative atomic masses : H=1, C=12) | |
| You must show your working. | |
| Deduce the molecular formula of this hydrocarbon. | |
| (d) A different hydrocarbon has a relative formula mass It has an empirical formula of CH ₂ . | of 84. |
| | |

DO NOT WRITE IN THIS AREA

AREA

DO NOT WRITE IN THIS

8 Calcium carbonate reacts with dilute hydrochloric acid to produce calcium chloride, water and carbon dioxide.

$$CaCO_3 + 2HCl \rightarrow CaCl_2 + H_2O + CO_2$$

(a) A student wanted to measure the amount of gas produced in two minutes.

The student suggested that this could be done by counting the number of bubbles formed.

However, the bubbles are produced too quickly to count them.

Figure 4 shows a conical flask in which the calcium carbonate and dilute hydrochloric acid are reacting.

Complete Figure 4 to show the apparatus that could be used to measure accurately the volume of gas given off in two minutes.

(2)




Figure 4

(b) The reaction between calcium carbonate and dilute hydrochloric acid is exothermic.

Explain, in terms of bond breaking and bond making, why some reactions are exothermic.

| | |
|------|--|
| | |
| | |

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

*(c) An investigation was carried out into the rate of reaction of calcium carbonate with dilute hydrochloric acid.

5.0g of small lumps of calcium carbonate were reacted with 50 cm³ of 0.50 mol dm⁻³ hydrochloric acid.

Another 5.0g of the same sized lumps of calcium carbonate were reacted with 50cm³ of 1.0 mol dm⁻³ hydrochloric acid.

The volume of gas collected in two minutes was recorded for each experiment.

The two experiments were then repeated, each using 5.0g of large lumps of calcium carbonate.

Figure 5 shows the results.

| concentration of | volume of gas collected in cm ³ | |
|--|--|----------------------------------|
| hydrochloric acid in mol dm ⁻³ | small lumps of calcium carbonate | large lumps of calcium carbonate |
| 0.50 | 17.2 | 3.1 |
| 1.0 | 35.1 | 5.6 |

Figure 5

Explain, in terms of collision of particles, how these results show the effect of the size of the lumps of calcium carbonate and the effect of the concentration of the acid on the rate of this reaction.

| (6) |
|-----|
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |

DO NOT WRITE IN THIS AREA

| (7.16.0 |
|---------------------------------------|
| (Total for Question 8 = 11 marks) |



DO NOT WRITE IN THIS AREA

| 9 | Fluorine, chlorine, bromine, iodine and astatine are elements in group 7. (a) Describe the test to show that a gas is chlorine. | (2) |
|---|--|-----|
| | (b) Bromine reacts with hydrogen to form hydrogen bromide. Hydrogen bromide dissolves in water to form a solution. State the name of the solution formed. | (1) |
| | (c) There is a trend in the colour and the state of the halogens at room temperature. Predict the colour and state of astatine at room temperature. colour | |

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(d) Bromine, chlorine and iodine are dissolved in water to make aqueous solutions. Potassium iodide solution is added to each of these solutions.

Figure 6 shows the observations.

| halogen | initial colour of aqueous solution | final colour of mixture |
|----------|------------------------------------|----------------------------|
| bromine | orange | brown |
| chlorine | pale green | brown |
| iodine | brown | brown |

Figure 6

| Explain the observations shown in the table. | (4) |
|--|-------------|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
|)Fluorine reacts vigorously with iron to produce iron(III) fluoride, FeF₃. | |
| Write the balanced equation for this reaction. | |
| | (2) |
| (Total for Question 9 = | = 11 marks) |
| (Total for Question 9 = | = 11 marks) |



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

10 (a) A sample of potassium carbonate is contaminated with a small amount of sodium carbonate.

When a flame test is carried out on the sample, a bright yellow flame is seen.

Describe how you could show that potassium and sodium ions are present in this sample.

(2)

(b) Hydrochloric acid reacts with a solution of sodium carbonate.

$$2HCl(aq) + Na_2CO_3(aq) \rightarrow 2NaCl(aq) + CO_2(g) + H_2O(l)$$

(3)

Write the ionic equation for this reaction.

(6)

DO NOT WRITE IN THIS AREA

*(c) A student tests solutions of three ionic substances, **K**, **L** and **M**.

The student carries out the same two tests on each of the three solutions.

Test 1 add dilute nitric acid and then silver nitrate solution.

Test 2 add a few drops of sodium hydroxide solution and warm the mixture.

Figure 7 shows the results of the tests and the student's conclusions about the identity of each substance.

| ionic substance | test 1 | test 2 | student's conclusion |
|-----------------|-------------------|---------------------|----------------------|
| K | white precipitate | colourless solution | ammonium chloride |
| L | white precipitate | white precipitate | aluminium chloride |
| М | no precipitate | green precipitate | iron(II) sulfate |

Figure 7

None of the student's conclusions are fully justified.

Explain which part of each conclusion is justified and what further work can be carried out to fully justify each conclusion.

| | |
|------|--|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |

DO NOT WRITE IN THIS AREA

| | (Total for Question 10 = 11 marks) | | |
|-----------------------------|------------------------------------|--|--|
| | | | |
| TOTAL FOR PAPER = 100 MARKS | | | |



The periodic table of the elements

| 9 | |
|-----|---|
| 2 | |
| 4 | |
| က | |
| | |
| | |
| | |
| | |
| | |
| | |
| | _ |
| | _ |
| | |
| | _ |
| | |
| | _ |
| | |
| | |
| 2 | |
| 1 2 | |

| _ | | | | | | |
|-----------------------|--|---------------------------------|------------------------------------|-------------------------------------|-------------------------------------|--------------------------------------|
| 0 | 4 He helium 2 | 20 Ne neon 10 | 40 Ar argon 18 | 84 Kr krypton 36 | 131 Xe xenon 54 | [222] Rn radon 86 |
| _ | | 19 fluorine 9 | 35.5 CI chlorine 17 | 80 Br bromine 35 | 127 | [210] At astatine 85 |
| 9 | | 16 O oxygen 8 | 32 S sulfur 16 | 79 Se selenium 34 | 128 Te tellunium 52 | [209] Po polonium 84 |
| 2 | | 14 N nitrogen 7 | 31 P phosphorus 15 | 75 As arsenic 33 | 122 Sb antimony 51 | 209 Bi bismuth 83 |
| 4 | | 12 C carbon 6 | 28 Si silicon 14 | 73 Ge germanium 32 | 119 Sn fin 50 | 207 Pb lead 82 |
| က | | 11 B boron 5 | 27 AI aluminium 13 | 70 Ga gallium 31 | 115 In indium 49 | 204 T thallium 81 |
| | | | | 65 Zn zinc 30 | 112 Cd cadmium 48 | 201 Hg mercury 80 |
| | | | | 63.5 Cu copper 29 | 108 Ag silver 47 | 197 Au gold 79 |
| | | | | 59 Ni nickel 28 | 106 Pd palladium 46 | 195 Pt platinum 78 |
| | | | | 59 Co cobalt 27 | 103 Rh rhodium 45 | 192 Ir iridium 77 |
| | 1 H hydrogen 1 | | | 56 Fe iron 26 | 101 Ru ruthenium 44 | 190 Os osmium 76 |
| • | | | | 55 Mn manganese 25 | [98] Tc technetium 43 | 186 Re rhenium 75 |
| | mass 30 umber | | 52 Cr chromium 24 | 96 Mo molybdenum 42 | 184 W tungsten 74 | |
| | Key relative atomic mass atomic symbol name atomic (proton) number | | | 51 V vanadium 23 | 93 Nb niobium 41 | 181 Ta tantalum 73 |
| relativ ato | | | | 48 Ti titanium 22 | 91 Zr zirconium 40 | 178 Hf hafnium 72 |
| | · | | | 45 Sc scandium 21 | 89 Y yttrium 39 | 139 La* Ianthanum 57 |
| 7 | | 9 Be beryllium | 24 Mg magnesium 12 | 40 Ca calcium 20 | 88 Sr strontium 38 | 137 Ba barum 56 |
| _ | | 7 Li Ilthium 3 | 23 Na sodium 11 | 39 K potassium 19 | 85 Rb rubidium 37 | 133 Cs caesium 55 |
| | | | | | | |

* The elements with atomic numbers from 58 to 71 are omitted from this part of the periodic table.

The relative atomic masses of copper and chlorine have not been rounded to the nearest whole number.

