



## Mark Scheme (Results)

January 2021

Pearson Edexcel International GCSE  
In Chemistry (4CH1) Paper 1C and Science  
(Double Award) (4SD0) Paper 1C

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

January 2021

Publications Code 4CH1\_1C\_2101\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2021

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks										
1 (a)	<table border="1" data-bbox="331 472 963 698"> <thead> <tr> <th data-bbox="331 472 651 510">Start</th> <th data-bbox="651 472 963 510">End</th> </tr> </thead> <tbody> <tr> <td data-bbox="331 510 651 548">solid</td> <td data-bbox="651 510 963 548">liquid</td> </tr> <tr> <td data-bbox="331 548 651 622">solid</td> <td data-bbox="651 548 963 622">gas</td> </tr> <tr> <td data-bbox="331 622 651 660">gas</td> <td data-bbox="651 622 963 660">liquid</td> </tr> <tr> <td data-bbox="331 660 651 698">liquid</td> <td data-bbox="651 660 963 698">gas</td> </tr> </tbody> </table>	Start	End	solid	liquid	solid	gas	gas	liquid	liquid	gas	<p data-bbox="991 443 1281 510">Award 1 mark for each correct row</p> <p data-bbox="991 551 1281 618"><b>ALLOW</b> gas to solid for sublimation</p>	3
Start	End												
solid	liquid												
solid	gas												
gas	liquid												
liquid	gas												
(b)	<p data-bbox="331 779 877 846">A description that refers to any three of the following points</p> <p data-bbox="331 887 938 920">M1 irregular /random arrangement (of particles)</p> <p data-bbox="331 960 919 1028">M2 large gaps between them /far apart /widely spaced</p> <p data-bbox="331 1068 794 1102">M3 random movement / move freely</p> <p data-bbox="331 1142 619 1176">M4 move (very) quickly</p>	<p data-bbox="991 960 1233 994"><b>ALLOW</b> spread out</p> <p data-bbox="991 1140 1273 1207"><b>IGNORE</b> references to kinetic energy</p>	3										
6 marks													

Question number	Answer	Notes	Marks
2 (a) (i)	A  A is the correct answer because 100°C is above the boiling point of W  B is not the correct answer because X is a solid at 100°C C is not the correct answer because Y is a solid at 100°C D is not the correct answer because Z is a solid at 100°C		1
(ii)	B  B is the correct answer because X is a liquid for 1840°C  A is not the correct answer because W is a liquid for 67°C C is not the correct answer because Y is a liquid for 1150°C D is not the correct answer because Z is a liquid for 330°C		1
(iii)	C  C is the correct answer because Y is a liquid at 1000°C and a gas at 2000°C  A is not the correct answer because W is a gas at 1000°C and at 2000°C B is not the correct answer because X is a liquid at 1000°C and 2000°C D is not the correct answer because Z is a gas at 1000°C and at 2000°C		1
(b)	ionic	<b>ALLOW</b> electrovalent	1
(c)	the (impure) substance will melt over a range of temperatures	<b>ALLOW</b> the (impure) substance will have a lower melting point	1
			5 marks

Question number	Answer	Notes	Marks
3 (a) (i)	M1 dissolving M2 diffusion	Answers can be in either order	2
(b) (i)	An explanation that links any two of the following points  M1 crystals dissolve faster  M2 (potassium iodide/ lead nitrate/ water) particles move faster / (lead/ iodide) ions move faster / rate of diffusion increases  M3 therefore (lead and iodide) ions/ particles meet / collide after a shorter period of time/ sooner	<b>ALLOW</b> (potassium iodide /lead nitrate/ water) particles have more energy  <b>ALLOW</b> molecules in place of particles if referring to water  <b>IGNORE</b> references to more collisions or more energetic collisions	2
(c) (i)	3 / three		1
(ii)	2+ /+2	<b>ALLOW</b> Pb <sup>2+</sup>	1
(d)	$\text{Pb}(\text{NO}_3)_2(\text{aq}) + 2\text{KI}(\text{aq}) \rightarrow \text{PbI}_2(\text{s}) + 2\text{KNO}_3(\text{aq})$	<b>ALLOW</b> multiples and fractions	1
			7 marks

Question number	Answer	Notes	Marks
4 (a)	<p>Example calculation</p> <p>M1 (volume of oxygen =) <math>100 - 25</math> <b>OR</b> <math>75</math> (cm<sup>3</sup>)</p> <p>M2 <math>75 \div 365 \times 100</math></p> <p>M3 20.5 (%)</p>	<p>Correct answer of 20.5 % with or without working scores 3</p> <p><b>ALLOW</b> ecf from M1</p> <p><b>ALLOW</b> ecf from M2</p> <p><b>ALLOW</b> 2 or more significant figures</p> <p><b>REJECT</b> incorrect rounding</p> <p>Use of 265 instead of 365 gives an answer of 28.3 and scores 2</p> <p>Alternative method</p> <p>M1 (volume of air left =) <math>265 + 25</math> <b>OR</b> <math>290</math> (cm<sup>3</sup>)</p> <p>M2 <math>290 \div 365 \times 100</math> <b>OR</b> <math>79.5</math> (%)</p> <p>M3 <math>(100 - 79.5 =)</math> 20.5 (%)</p>	3
(b) (i)	<p>M1 paint provides a barrier</p> <p>M2 which prevents oxygen / water getting to /reacting with the iron</p>	<p><b>ALLOW</b> paint forms a coating (on the iron) / paint is non-permeable</p> <p><b>ALLOW</b> air</p>	2
(b) (ii)	<p>M1 zinc is more reactive/higher in the reactivity series (than iron)</p> <p>M2 zinc will oxidise / react / corrode instead of /before iron</p>	<p><b>ALLOW</b> zinc is a sacrificial metal</p> <p><b>IGNORE</b> references to zinc rusting</p> <p><b>IGNORE</b> references to galvanising</p>	2
			7 marks

Question number	Answer	Notes	Marks					
5 (a)	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th data-bbox="328 300 971 338" style="text-align: center;">Method</th> </tr> </thead> <tbody> <tr> <td data-bbox="328 338 971 376">filtration</td> </tr> <tr> <td data-bbox="328 376 971 414">simple distillation <b>or</b> fractional distillation</td> </tr> <tr> <td data-bbox="328 414 971 524">fractional distillation</td> </tr> <tr> <td data-bbox="328 524 971 562">crystallisation</td> </tr> </tbody> </table>	Method	filtration	simple distillation <b>or</b> fractional distillation	fractional distillation	crystallisation	<p><b>ALLOW</b> filtering  <b>ALLOW</b> distillation  <b>REJECT</b> simple distillation or distillation</p>	4
Method								
filtration								
simple distillation <b>or</b> fractional distillation								
fractional distillation								
crystallisation								
(b) (i)	<p>M1 A and B</p> <p>M2 because they are the same height /moved the same distance up the paper / have the same <math>R_f</math> values as the spots in the purple ink</p>	M2 dep on M1 correct or missing	2					
(b) (ii)	<p>M1 D</p> <p>M2 because the spot is closest to the start line /travelled the least distance (from the start line) / has the lowest <math>R_f</math> value</p>	M2 dep on M1 correct or missing	2					
(c)	<p>Example calculation</p> <p>M1 <math>120 \times 0.72</math></p> <p>M2 <math>86 / 86.4(\text{mm})</math></p>	Correct answer of 86 or 86.4 (mm) with or without working scores 2	2					
			10 marks					



Question number	Answer	Notes	Marks						
6 (a)	<table border="1"> <tr> <td data-bbox="327 309 646 380">precipitate of barium carbonate</td> <td data-bbox="647 309 968 380">precipitate of barium sulfate</td> </tr> <tr> <td data-bbox="327 383 646 454">no precipitate</td> <td data-bbox="647 383 968 454">no precipitate</td> </tr> <tr> <td data-bbox="327 456 646 528">precipitate of calcium carbonate</td> <td data-bbox="647 456 968 528">precipitate of calcium sulfate</td> </tr> </table>	precipitate of barium carbonate	precipitate of barium sulfate	no precipitate	no precipitate	precipitate of calcium carbonate	precipitate of calcium sulfate	<p>if barium sulfate and calcium carbonate correct but without including 'precipitate of' scores 1 out of 2</p> <p><b>ALLOW</b> correct formulae</p>	3
precipitate of barium carbonate	precipitate of barium sulfate								
no precipitate	no precipitate								
precipitate of calcium carbonate	precipitate of calcium sulfate								
(b)	<p>A description that refers to any six of the following points</p> <p>M1 do a flame test</p> <p>M2 sodium chloride produces a yellow flame</p> <p>M3 add acid</p> <p>M4 potassium carbonate effervesces / bubbles</p> <p>M5 add dilute nitric acid</p> <p>M6 add silver nitrate (solution)</p> <p>M7 potassium chloride gives a white precipitate</p> <p>M8 potassium iodide gives a yellow precipitate</p>	<p><b>ACCEPT</b> any description of a flame test</p> <p><b>ACCEPT</b> yellow-orange or orange</p> <p><b>IGNORE</b> any flame colour given for the potassium compounds</p> <p><b>ALLOW</b> any named acid</p> <p><b>ACCEPT</b> carbon dioxide/gas given off which turns limewater cloudy for M4</p> <p>M4 is dep on M3</p> <p>M7 and M8 are dep on M6</p> <p><b>ALLOW</b> addition of chlorine/bromine solution as an alternative to M6</p> <p>M7 no colour change with potassium chloride</p>	6						

M8 solution turns brown  
with potassium iodide

If this alternative given no  
M5

9 marks

Question number	Answer	Notes	Marks
7 (a)	<p>M1 two lithium atoms each lose one electron /give one electron to oxygen</p> <p>M2 oxygen gains two electrons</p> <p>M3 lithium (ion) has an electron configuration of 2 and oxide (ion) is 2,8</p>	<p><b>ALLOW</b> lithium loses one electron /gives one electron to oxygen</p> <p><b>ALLOW</b> oxygen becomes 2,8</p> <p>All 3 marks can be scored from diagrams showing the electron configurations of the ions</p> <p>0 marks if reference to sharing electrons</p>	3
(b) (i)	<p>M1 (temperature after) = 27.7 °C</p> <p>M2 temperature rise = 10.4 °C</p>	<p><b>ALLOW</b> ecf from M1</p>	2
(ii)	<p>Example calculation</p> <p>M1 Use of 100 in <math>Q = m \times c \times \Delta T</math></p> <p>M2 Use of 10.4 in <math>Q = (m \times) c \times \Delta T</math></p> <p>M3 4368J</p> <p>M4 4400J</p>	<p>Correct answer of 4400J with or without working scores 4</p> <p><b>ALLOW</b> ecf from (b)(i)</p> <p>100 x 4.2 x 10.4 scores M1 and M2</p> <p><b>ALLOW</b> ecf from M1 and M2</p> <p><b>ALLOW</b> ecf from M3</p>	4
(iii)	<p>Example calculation</p> <p>M1 <math>5210 \div 1000</math> or 5.21</p> <p>M2 <math>5.21 \div 0.0580</math></p> <p>M3 -89.8(kJ/mol)</p>	<p><b>IGNORE</b> + or - sign in front of answer</p> <p>Correct answer of -89.8 (kJ/mol) scores 3</p> <p><b>ALLOW</b> -90 (kJ/mol) or any number of sig figs as long as correctly rounded.</p>	3
(iv)	<p>polystyrene is a good insulator /poor conductor (of heat) OR to minimise/reduce heat loss</p>	<p><b>ALLOW</b> prevent heat loss</p>	1
			13 marks

Question number	Answer	Notes	Marks
8 (a)	M1 solid  M2 dark grey / black		2
(b) (i)	Example calculation  M1 $(51 \times 79) + (49 \times 81)$ <b>OR</b> 7998  M2 $7998 \div 100$  M3 80.0	80.0 with no working scores 3      79.9 with no working scores 1   79.98 or 80 with no working scores 2	3
	(ii) same electron configuration	<b>ALLOW</b> same (total) number of electrons  <b>IGNORE</b> same number of electrons in the outer shell  <b>IGNORE</b> references to same number of protons	1
(c) (i)	An explanation that links the following three points  M1 the order of reactivity is chlorine (most), bromine and iodine (least)  M2 chlorine (is most reactive because it) displaces bromine and iodine/ oxidises bromide and iodide (ions) / reacts with sodium bromide and sodium iodide  M3 bromine (is less reactive than chlorine since it) only displaces iodine / only oxidises iodide (ions) / only reacts with sodium iodide	<b>ACCEPT</b> bromine is only displaced by chlorine and iodine is displaced by chlorine and bromine scores M2 and M3  <b>ALLOW</b> chlorine has two reactions, bromine has one reaction and iodine no reactions for 1 mark out of M2 and M3  Deduct 1 mark for incorrect use of ine	3

		and ide e.g. bromine displaces iodide	
(ii)	bromine cannot displace itself / bromine does not react with sodium bromide OWTTE	<b>ALLOW</b> there would be no reaction	1 2
(iii)	M1 bromine is reduced and iodide (ions)/I <sup>-</sup> is oxidised M2 bromine gains electrons and iodide (ions)/I <sup>-</sup> loses electrons  <b>OR</b> M1 bromine gains electrons so is reduced M2 iodide (ions) /I <sup>-</sup> loses electrons so is oxidised	Deduct 1 mark for mention of iodine (ions) being oxidised or losing electrons  <b>REJECT</b> iodine (ions) loses electrons so is oxidised	

12 marks

Question number	Answer	Notes	Marks
9 (a)	M1 (propane/it) contains hydrogen and carbon (atoms)  M2 only	<b>REJECT</b> carbon and hydrogen molecules  M2 is dependent on mention of just carbon and hydrogen in M1	2
(b) (i)	carbon monoxide	<b>ALLOW</b> CO	1
(ii)	it decreases the capacity of the blood to transport oxygen OWTTE	<b>ALLOW</b> carbon monoxide binds to haemoglobin	1
(c)	M1 (strong electrostatic) attraction between (bonding) pair of electrons  M2 (and) nuclei (of both atoms)  <b>OR</b>  M1 (bonding) pair of electrons  M2 attracted to nuclei	<b>REJECT</b> nucleus    <b>REJECT</b> nucleus  0 marks if reference to intermolecular forces between atoms	2
(d)	An explanation that links the following three points  M1 (crude oil) produces more long chain hydrocarbons than can be used directly OWTTE  M2 shorter chain alkanes are more flammable /more useful as fuels  M3 alkenes are used to make polymers / plastics	<b>ALLOW</b> less demand for long chain hydrocarbons  <b>ALLOW</b> shorter chain alkanes/hydrocarbons are more useful	3
(e) (i)	M1 C <sub>3</sub> H <sub>7</sub> Br M2 HBr	<b>ALLOW</b> polysubstituted product if correct balancing number in front of Br <sub>2</sub> and HBr	2
(ii)	substitution		1

Question number	Answer	Notes	Marks
10 (a) (i)	curve of best fit	<b>REJECT</b> dot to dot line	1
(ii)	M1 lines shown on graph	<b>ALLOW</b> extra point on curve at 7 carbon atoms	2
(iii)	M2 value correctly read from graph (expected value between 97 and 103°C)	<b>ACCEPT</b> value to $\pm 1^\circ\text{C}$	3
	An explanation that links the following three points		
	M1 the boiling point increases as the number of carbons / the chain length increases	<b>ALLOW</b> boiling point increases as the $M_r$ increases	
	M2 because the intermolecular forces (of attraction) get stronger	<b>REJECT</b> directly proportional	
	M3 and therefore take more energy to overcome / break	M3 dep on M2 Any mention of breaking covalent bonds does not score M2 or M3	
(b)	M1 same <b>molecular</b> formula		2
	M2 different <b>displayed</b> / <b>structural</b> formulae	<b>ALLOW</b> different structures/ different arrangement of atoms	
(c) (i)	M1 $82.8 \div 12$ <b>or</b> 6.9 $17.2 \div 1$ <b>or</b> 17.2	0 marks if upside down calculation or use of atomic numbers	2
	M2 (divide by smallest to give) 1:2.5 which is 2:5	<b>ACCEPT</b> alternative methods	
(ii)	$\text{C}_4\text{H}_{10}$		1

(d)	M1 moles of CO <sub>2</sub> = 7 <b>or</b> X = 7 M2 moles of H <sub>2</sub> O = 8 <b>or</b> Y = 8 M3 balancing number = 11 <b>or</b> Z = 11	<b>ALLOW</b> ecf from incorrect values of X and Y	3
			14 marks

Question number	Answer	Notes	Marks
11 (a) (i)	glowing splint relights	<b>REJECT</b> burning splint	1
(ii)	A description that refers to the following three points M1 filter out manganese(IV) oxide / solid M2 leave to dry M3 same mass/ 1g of manganese(IV) oxide / solid		3
(b) (i)	M1 280 ÷ 120 M2 2.33	<b>ALLOW</b> ecf from M1 <b>ALLOW</b> any number of significant figures except 1	2
(ii)	An explanation that links the following three points M1 the concentration of hydrochloric acid is greatest M2 therefore there are more collisions M3 per unit time	<b>ALLOW</b> the surface area of zinc is greatest <b>ALLOW</b> greatest number of/more particles (of hydrochloric acid/ zinc) More frequent collisions scores M2 and M3	3 2
(iii)	M1 curve above original and starts at 0 M2 curve goes flat at same volume (410cm <sup>3</sup> )	Max 1 if incorrect reference to energy	



(iv)	<p>M1 greater surface area</p> <p>M2 more collisions per unit time / more frequent collisions</p>		2
(c)	<p>M1 <math>8.46 \times 10^{-3}</math> mol of zinc</p> <p>M2 therefore <math>1.69 \times 10^{-2}</math> mol hydrochloric acid needed (which is less than <math>2.50 \times 10^{-2}</math> mol)</p> <p><b>OR</b></p> <p>M1 <math>1.25 \times 10^{-2}</math> mol of zinc are needed</p> <p>M2 therefore 0.8(13) g of zinc is needed (and there is only 0.55g)</p>	<p><b>ALLOW</b> any number of sig figs including one e.g. 0.008 moles of zinc, therefore 0.016 moles of acid needed scores M1 and M2</p>	2
			15 marks

